

References

Chapter 1: The Power of Positive Thinking

For the Mayo Clinic study of optimists and pessimists, see Toshihiko Maruta MD et al., Mayo Clinic Proceedings, August 2002, or visit: <http://www.sciencedaily.com/releases/2002/08/020813071621.htm>

For the 2004 study involving 999 Dutch men, see E.J. Giltay et al., 'Dispositional optimism and all-cause and cardiovascular mortality in a prospective cohort of elderly Dutch men and women', *Archives of General Psychiatry*, 2004, 61, 1126–35.

For the nun study, see D.D. Danner et al., 'Positive emotions in early life and longevity: findings from the nun study', *Journal of Personality and Social Psychology*, 2001, 80(5), 804–13.

For the study where volunteers were exposed to the cold or influenza viruses, see S. Cohen et al., 'Positive emotional style predicts resistance to illness after experimental exposure to rhinovirus or influenza A virus', *Psychosomatic Medicine*, 2006, 68, 809–15.

For an article containing the study of 200 telecommunications executives, see Peggy Rynk, 'The value of a healthy attitude: how faith, anger, humor, and boredom can affect your health', *Vibrant Life*, March–April 2003.

For the study of 586 people finding that attitude is the best prevention against heart disease, see D.M. Becker, 'Positive attitude is best prevention against heart disease', paper presented at the American Heart Association Annual Scientific Sessions, Anaheim, CA, 12 November 2001. To read an account of the study, visit: http://www.hopkinshospital.org/health_info/Heart/Reading/positive_attitude

For the study of 866 heart patients and positive attitude, see B. Brummett, 'Positive outlook linked to longer life in heart patients', paper presented at the American Psychosomatic Society, March 2003. To read an account of the study, visit: <http://dukemednews.org/av/medminute.php?id=6511>

For the emotional vitality study, see L.D. Kubzansky and R.C. Thurston, 'Emotional vitality and incident coronary heart disease: benefits of healthy psychological functioning', *Archives of General Psychiatry*, 2007, 64(12), 1393–1401.

For the 'Hard Marriage, Hard Heart' research, see:

- T.W. Smith et al., 'Marital conflict behavior and coronary artery calcification', paper presented at the American Psychosomatic Society 64th Annual Meeting, Denver, CO, 3 March 2006.
- T.W. Smith et al., 'Hostile personality traits and coronary artery calcification in middle-aged and older married couples: different effects for self-reports versus spouse ratings', *Psychosomatic Medicine*, 2007, 69(5), 441–48.

- P. Pearsall, 'Contextual cardiology: what modern medicine can learn from ancient Hawaiian wisdom', *Cleveland Clinical Journal of Medicine*, 2007, 74(1), S99–S104.

For the 25-year hostility study, see J.C. Barefoot et al., 'Hostility, CHD incidence, and total mortality: A 25-year follow-up study of 255 physicians', *Psychosomatic Medicine*, 1983, 45(1), 59–63.

For the paper quoting hostility as an indicator of heart disease, see R.B. Williams et al., 'Psychosocial risk factors for cardiovascular disease: more than one culprit at work', *Journal of the American Medical Association*, 2003, 290(16), 2190–92.

For the Finnish satisfaction study, see H. Koivumaa-Honkanen et al., 'Self-reported life satisfaction and 20-year mortality in healthy Finnish adults', *American Journal of Epidemiology*, 2000, 152(10), 983–91.

For the 'Money buys happiness' research, see Elizabeth Dunn, 'Money buys happiness when you spend on others: UBC and Harvard research', University of British Columbia Media Release, 20 March 2008. To view the media release, visit: <http://www.publicaffairs.ubc.ca/media/releases/2008/mr-08-032.html>

For the effect of attitude on aging, see B.R. Levy et al., 'Longevity increased by positive self-perceptions of aging', *Journal of Personality and Social Psychology*, 2002, 83(2), 261–70.

For the study of positive attitude being good for blood pressure, see G.V. Ostir et al., 'Hypertension in older adults and the role of positive emotions', *Psychosomatic Medicine*, 2006, 68, 727–33.

For the study linking frailty with attitude, see G.V. Ostir et al., 'Onset of frailty in older adults and the protective role of positive affect', *Psychology and Aging*, 2004, 19(3), 402–08.

For the link between life satisfaction and longevity, see T.M. Lyyra et al., 'Satisfaction with present life predicts survival in octogenarians', *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 2006, 61, 319–26.

For the Posit Science Corporation study, see:

- H.W. Mahncke et al., 'Memory enhancement in healthy older adults using a brain plasticity-based training program: a randomized, controlled study', *Proceedings of the National Academy of Sciences, USA*, 2006, 103(33), 12523–28.
- H.W. Mahncke et al., 'Brain plasticity and functional losses in the aged: scientific basis for a novel intervention', *Progress in Brain Research*, 2006, 157, 81–109.

For the Harvard age study going back to 1959, see Ellen J. Langer PhD, *Mindfulness* (Da Capo Press, 1990).

For information on using the brain and reducing the risk of Alzheimer's, see R.S. Wilson et al., 'Participation in cognitively stimulating activities and risk of incident Alzheimer's disease', *Journal of the American Medical Association*, 2002, 287(6), 742–48.

For the priming experiments, see:

- J.A. Bargh et al., 'Automaticity of social behavior: direct effects of trait construct and stereotype activation on action', *Journal of Personality and Social Psychology*, 1996, 71(2), 230–44.
- T.M. Hess et al., 'Explicit and implicit stereotype activation effects on memory: do age and awareness moderate the impact of priming?', *Psychology and Aging*, 2004, 19(3), 495–505.

Chapter 2: The Power of Believing

The quotation from Professor Benedetti can be found in F. Benedetti, 'Mechanisms of placebo and placebo-related effects

across diseases and treatments', *Annual Review of Pharmacology and Toxicology*, 2008, 46, 33–60. This is also a good review of recent research into the placebo effect.

For the release of dopamine when Parkinson's patients receive placebos, see:

- R. de la Fuente-Fernández et al., 'Expectation and dopamine release: mechanism of the placebo effect in Parkinson's disease', *Science*, 2001, 293(5532), 1164–66.
- R. de la Fuente-Fernández et al., 'Dopamine release in human ventral striatum and expectation of reward', *Behavioral Brain Research*, 2002, 136(2), 359–63.

The first evidence of opioid release during placebo analgesia can be found in J.D. Levine et al., 'The mechanism of placebo analgesia', *Lancet*, 1978, 654–57.

For an account of placebo effects tracking closely the treatments with which they're paired, and a discussion of the similarity in brain scans while patients receive Prozac or a placebo, see F. Benedetti et al., 'Neurobiological mechanisms of the placebo effect', *Journal of Neuroscience*, 2005, 25(45), 10390–402.

For the MRI brain scans of people receiving placebos and high placebo responses, see:

- J.K. Zubieta et al., 'Placebo effects mediated by endogenous opioid activity on m-opioid receptors', *Journal of Neuroscience*, 2005, 25, 7754–62.
- T.D. Wager et al., 'Placebo effects on human m-opioid activity during pain', *Proceedings of the National Academy of Sciences, USA*, 2007, 104(26), 11056–61.

For some clinical trial results, visit: <https://www.centerwatch.com/cwweekly/category/news/trial-results/> (last accessed, 5th June 2018).

For the chronic fatigue syndrome study that tested acyclovir, see S.E. Strauss et al., 'Acyclovir treatment of the chronic fatigue syndrome. Lack of efficacy in a placebo-controlled trial', *New England Journal of Medicine*, 1988, 319(26), 1692–98.

For the placebo studies on asthma, see:

- T. Luparello et al., 'Influences of suggestion on airway reactivity in asthmatic subjects', *Psychosomatic Medicine*, 1969, XXX, 819–25.
- E.R. McFadden et al., 'The mechanism of action of suggestion in the induction of acute asthma attacks', *Psychosomatic Medicine*, 1969, XXXI, 134–43.

For the study of performance-enhancing placebos, see F. Benedetti et al., 'Opioid-mediated placebo responses boost pain endurance and physical performance: is it doping in sport competitions?' *Journal of Neuroscience*, 2007, 27(44), 11934–39.

For the study involving the hotel-room maids, see A. J. Crum and E. J. Langer, *Psychological Science*, 2007, 18(2), 165–171.

For the study of women's beliefs affecting their mathematics performance, see I. DarNimrod and S.J. Heine, 'Exposure to scientific theories affects women's math performance', *Science*, 2006, 314(5798), 435.

For information on placebo responders, as well as lots of information on a range of placebo studies, see Daniel Moerman, *Meaning, Medicine and the 'Placebo Effect'* (Cambridge University Press, 2002).

The 1954 bleeding ulcers study was reported in Daniel Moerman's book.

For the study involving hidden arm pain, see F. Benedetti, 'The opposite effects of the opiate antagonist naloxone and the

cholecystokinin antagonist proglumide on placebo analgesia', *Pain*, 1996, 64(3), 535–43.

For the dental injections given with an 'oversell' or 'undersell' message, see S.L. Gryll and M. Katahn, 'Situational factors contributing to the placebo effect', *Psychopharmacology*, 1978, 57(3), 253–61.

For the study on the effect of positive and negative consultations, see K.B. Thomas, 'General practice consultations: is there any point in being positive?' *British Medical Journal*, 1987, 294, 1200–02.

For the studies of optimists and pessimists and how they respond in the placebo effect, see the following:

- For the pessimists study, see A.L. Geers et al., 'Reconsidering the role of personality in placebo effects: dispositional optimism, situational expectations, and the placebo response', *Journal of Psychosomatic Research*, 2005, 58(2), 212–17.
- For the optimists study, see A.L. Geers et al., 'Further evidence for individual differences in placebo responding: an interactionist perspective', *Psychosomatic Research*, 2007, 62(5), 563–70.

For the conditioned immunosuppression using cyclosporin A, see M. U. Goebel et al., 'Behavioral conditioning of immunosuppression is possible in humans', *FASEB Journal*, 2002, 16, 1869–73.

For Benedetti's conditioning of immune and growth hormone levels, see F. Benedetti et al., 'Conscious expectation and unconscious conditioning in analgesic, motor, and hormonal placebo/nocebo responses', *Journal of Neuroscience*, 2002, 23, 4315–23.

For the Benedetti study where Parkinson's patients received placebos instead of apomorphine, see F. Benedetti et al., 'Teaching Neurons to Respond to Placebos', *Journal of Physiology*, 2016, 594(19), 5647-5660

For PCDR studies and a discussion on how to use conditioning to boost the placebo effect, see Jo Marchant, *Cure* (Cannongate Books, London, 2017), chapter 3, 'Pavlov's Power'.

Chapter 3: Drugs Work Better If We Believe in Them

For the 2008 antidepressants analysis that found over 80 per cent placebo effect, see I. Kirsch et al., 'Initial severity and antidepressant benefits: a meta-analysis of data submitted to the food and drug administration', *PLoS Medicine*, February 2008, 5(2), e45, 0260–68.

A number of the studies reported in this chapter were also cited in Daniel Moerman's excellent book, cited above. I thoroughly recommend it to anyone seeking a fuller understanding of the placebo effect and its implications.

For the quote from C.G. Helman, see his chapter entitled 'Placebos and nocebos: the cultural construction of belief' in *Understanding the Placebo Effect in Complementary Medicine, Theory, Practice and Research*, ed. D. Peters (Churchill Livingstone, 2001).

For the study of blue and pink sedatives and stimulants, see B. Blackwell et al., 'Demonstration to medical students of placebo responses and non-drug factors', *Lancet*, 1972, 1(7763), 1279–82.

For the study of injections versus tablets in the USA and Europe, see A.J. de Craen et al., 'Placebo effect in the acute treatment of migraine: subcutaneous placebos are better than oral placebos', *Journal of Neurology*, 2000, 247(3), 183–88.

For the study of Tagamet performed in France, see R. Lambert et al., 'Treatment of duodenal and gastric ulcer with cimetidine. A multicenter double-blind trial', *Gastroenterologie Clinique et Biologique*, 1977, 1(11), 855–60.

For the study in Brazil, see J.A. Salgado et al., 'Endoscopic findings after antacid, cimetidine and placebo for peptic ulcer – importance of staging the lesions', *Arquivos De Gastroenterologia*, 1981, 18(2), 51–3. Both studies are reported in Daniel Moerman's book, cited above.

The reduction in effectiveness of Tagamet once Zantac was available is described in Daniel Moerman's book. It's also mentioned in Herbert Benson MD's *Timeless Healing* (Scribner, 1995).

For the University of Keel aspirin study, see A. Branthwaite and P. Cooper, 'Analgesic effects of branding in treatment of headaches', *British Medical Journal*, 1981, 282, 1576–78.

For the suggestion that Viagra is enhanced by its name, see A.K. Vallance, 'Something out of nothing: the placebo effect', *Advances in Psychiatric Treatment*, 2006, 12, 287–96.

For the study of four placebos being better than two in anti-ulcer trials, see A.J. de Craen et al., 'Placebo effect in the treatment of duodenal ulcer', *British Journal of Clinical Pharmacology*, 1999, 48(6), 853–60.

For adherence in the clofibrate trial, see Coronary Drug Project Research Group, 'Influence of adherence to treatment and response of cholesterol on mortality in the coronary drug project', *New England Journal of Medicine*, 1980, 303(18), 1038–41.

For the real versus sham mammary ligation studies, see:

- E.G. Dimond et al., 'Comparison of internal mammary artery ligation and sham operation for angina pectoris', *American Journal of Cardiology*, 1960, 5, 483–86.

- L.A. Cobb et al., 'An evaluation of internal-mammary-artery ligation by a double-blind technique', *New England Journal of Medicine*, 1959, 260(22), 1115–18.

For the summary analysis of 53 trials comparing actual surgery with sham surgery, see K. Wartolowska et al., 'Use of placebo controls in the evaluation of surgery: Systematic review', *British Medical Journal*, 2014, 348, g3253.

For the naproxen study where patients either knew or didn't know that they were receiving it, see J.F. Bergmann et al., 'A randomised clinical trial of the effect of informed consent on the analgesic activity of placebo and naproxen in cancer pain', *Clinical Trials Meta-Analysis*, 1994, 29(1), 41–7.

For research showing that a placebo can be as good as 6–8mg of morphine, see J.D. Levine and N.C. Gordon, 'Influence of the method of drug administration on analgesic response', *Nature*, 1984, 312, 755.

For research showing that diazepam doesn't work unless a person knows they're receiving it, see F. Benedetti et al., 'Hidden administration of drugs', *Clinical Pharmacology & Therapeutics*, 2011, 90, 651–661.

The Benedetti quote, 'The existence of the placebo effect suggests that we must broaden our conception of the limits of... human capability,' can be found in F. Benedetti et al., 'Neurobiological mechanisms of the placebo effect', *Journal of Neuroscience*, 2005, 25(45), 10390–402.

For research showing that open label placebos work through nonconscious processes, see: K.B. Jensen et al., 'A neural mechanism for nonconscious activation of conditioned placebo and nocebo responses', *Cerebral Cortex*, 2015, 25, 3903–3910.

For the OLP study on IBS, see: T.J. Kaptchuk et al., 'Placebos Without Deception: A randomized controlled trial in irritable bowel syndrome', *PLoS ONE*, 2010, 5(12), e15591.

For the OLP study on chronic low back pain, see: C. Carvalho et al., 'Open-label Placebo Treatment in Chronic Low Back Pain: A randomized controlled trial', *Pain*, 2016, 157(12), 2766 – 2772.

For the OLP study on fatigue suffered by cancer patients, see: T.W. Hoenemeyer et al., 'Open-Label Placebo Treatment for Cancer-Related Fatigue: A Randomized-Controlled Clinical Trial', *Scientific Reports*, 2018, 8(2784), 1–8.

Chapter 4: The Power of Plasticity

For the study of brain changes in symphony musicians, see V. Sluming et al., 'Voxel-based morphometry reveals increased gray matter density in Broca's area in male symphony orchestra musicians', *NeuroImage*, 2002, 17(3), 1613–22.

The reference to brain-map changes as blind people learn Braille can be found in Norman Doidge MD, *The Brain That Changes Itself* (Penguin, 2007).

For the study of brain changes while students were studying for exams, see B. Draganski et al., 'Temporal and spatial dynamics of brain structure changes during extensive learning', *Journal of Neuroscience*, 2006, 26(23), 6314–17.

For the London taxi drivers study, see E.A. Maguire et al., 'London taxi drivers and bus drivers: a structural MRI and neuropsychological analysis', *Hippocampus*, 2006, 16, 1091–101.

For the study showing brain changes in mathematicians, see K. Aydin et al., 'Increased gray matter density in the parietal cortex

of mathematicians: a voxel-based morphometry study', *American Journal of Neuroradiology*, 2007, 28(10), 1859–64.

For the study of the effects of meditation on the brain, see S.W. Lazar et al., 'Meditation experience is associated with increased cortical thickness', *Neuroreport*, 2005, 16(17), 1893–97.

The Eric Kandel quote, as well as a discussion of brain changes due to psychotherapy, can be found in Norman Doidge's book, cited above.

For the study showing 15 per cent increase in hippocampal volume due to environmental enrichment, see G. Kemperman et al., 'Neuroplasticity in old age: sustained fivefold induction of hippocampal neurogenesis by long-term environmental enrichment', *Annals of Neurology*, 2002, 52, 135–143. See also L. Lu et al., 'Modification of hippocampal neurogenesis and neuroplasticity by social environments', *Experimental Neurology*, 2003, 183(2), 600–09.

For discovery of neurogenesis in the hippocampus, see P.S. Eriksson et al., 'Neurogenesis in the adult human hippocampus', *Nature Medicine*, 1998, 4(11), 1313–17.

The identification of neurogenesis right up to our final days can be found in the paper cited immediately above, on the discovery of neurogenesis in adults. The scientists received permission from terminally ill patients to inject them with a biological marker, bromodeoxyuridine (BrdU). When the patients died, neurogenesis was discovered.

For a good review of neurogenesis, see P. Taupin and F.H. Gage, 'Adult neurogenesis and neural stem cells of the central nervous system in mammals', *Journal of Neuroscience Research*, 2002, 69, 745–49.

Chapter 5: The Mind Can Heal the Body

For a good description of the brain and how it changes with our thoughts and emotions, see Joe Dispenza, *Evolve Your Brain* (Health Communications Inc., 2007).

For the research showing that hostility slows the rate of wound healing, see J.K. Kiecolt-Glaser et al., 'Hostile marital interactions, proinflammatory cytokine production, and wound healing', *Archives of General Psychiatry*, 2005, 62(12), 1377–84.

For the effects of stress on growth hormone levels and on the 'upregulation' and 'downregulation' of genes at wound sites, see S. Roy, et al., 'Wound site neutrophil transcriptome in response to psychological stress in young men', *Gene Expression*, 2005, 12(4–6), 273–87.

For social support speeding up wound healing, see C.E. Detillion et al., 'Social facilitation of wound healing', *Psychoneuroendocrinology*, 2004, 29(8), 1004–11.

Trumping our genes refers to the science of epigenetics. To learn more about epigenetics, see Bruce Lipton, *The Biology of Belief* (Mountain of Love/Elite Books, 2005). See also Dawson Church, *The Genie in Your Genes* (Elite Books, 2007).

For research on the effect of love and care on the growth of children, see David R. Hamilton PhD, *Why Kindness is Good for You* (Hay House, London, 2010), Chapter 10, 'Why Babies Need Love'. See also: www.bucharestearlyinterventionproject.org (last accessed 23 May 2018).

For information on neurogenesis and the suggestion that the mind interacts with DNA in stem cells, see Ernest L. Rossi, *The Psychobiology of Gene Expression* (Norton, 2002).

Chapter 6: The Power of Imagination and Observation

For the Benedetti study involving capsaicin, see F. Benedetti et al., 'Somatotopic activation of opioid systems by target-directed expectations of analgesia', *The Journal of Neuroscience*, 1999, 19(9), 3639–48.

For the study where experimental pain was induced in fingers, see G. Montgomery and I. Kirsch, 'Mechanisms of placebo pain reduction: an empirical investigation', *Psychological Science*, 1996, 7(3), 174–76.

In an email communication, I asked Fabrizio Benedetti whether a person with two different ailments would get better from one of them if he or she was given a placebo for that specific ailment (believing it to be a real drug) but not the other. Would a person with Parkinson's, say, who also had a headache and who received an anti-Parkinson's drug (which was really a placebo) find the tremors reducing but not the headache, and vice versa? Benedetti agreed that this would most likely be the case.

For the Karolinska Institute research showing that imagined movements of fingers, toes and tongue activated brain regions that governed them, see H. H. Ehrsson et al., 'Imagery of voluntary movement of fingers, toes, and tongue activates corresponding body-part-specific motor representations', *Journal of Neurophysiology*, 2003, 90(5), 3304–16.

For the piano study, see A. Pascual-Leone et al., 'Modulation of muscle responses evoked by transcranial magnetic stimulation during the acquisition of new fine motor skills', *Journal of Neurophysiology*, 1995, 74(3), 1037–45.

For the study where volunteers' fingers got 35 per cent stronger through imagined training, see V.K. Ranganathan et al., 'From mental power to muscle power – gaining strength by using the

mind', *Neuropsychologia*, 2004, 42(7), 944–56. See also G. Yue and K.J. Cole, 'Strength increases from the motor program: comparison of training with maximal voluntary and imagined muscle contractions', *Journal of Neurophysiology*, 1992, 67(5), 1114–23.

For the study showing the differences in muscle activation depending upon the imagined weight, see A. Guillot et al., 'Muscular responses during motor imagery as a function of muscle contraction types', *International Journal of Psychophysiology*, 2007, 66(1), 18–27.

For the study where the tetraplegic person opened an email with his mind, see L.R. Hochberg et al., 'Neuronal ensemble control of prosthetic devices by a human with tetraplegia', *Nature*, 2006, 442, 164–71.

For research into mental walking in virtual-reality simulators, see G. Pfurtscheller, et al., 'Walking from thought', *Brain Research*, 2006, 1071(1), 145–52.

For mirror neurons research where volunteers watched hands, mouth or foot movements, see G. Buccino et al., 'Action observation activates premotor and parietal areas in a somatotopic manner: an fMRI study', *European Journal of Neuroscience*, 2001, 13(2), 400–04.

For the 'Bend it like Beckham' paper, see P. Bach and S.P Tipper, 'Bend it like Beckham: embodying the motor skills of famous athletes', *Quarterly Journal of Experimental Psychology*, 2006, 59(12), 2033–39.

For the study where volunteers increased finger strength through watching training, see C.A. Porro et al., 'Enhancement of force after action observation: behavioural and neurophysiological studies', *Neuropsychologia*, 2007, 45(13), 3114–21.

For improvements gained by stroke patients through watching people perform routine actions, see D. Ertelt et al., 'Action observation has a positive impact on rehabilitation of motor deficits after stroke', *NeuroImage*, 2007, 36, Supplement 2, T164–73.

For activation of mirror neurons while watching someone playing a guitar, see G. Buccino et al., 'Neural circuits underlying imitation learning of hand actions: an event-related fMRI study', *Neuron*, 2004, 42, 323–34.

For activation of the brain by listening to sentences describing motion, see G. Buccino et al., 'Listening to action-related sentences modulates the activity of the motor system: a combined TMS and behavioral study', *Cognitive Brain Research*, 2005, 24(3), 355–63.

See also M. Tettamanti et al., 'Listening to action-related sentences activates fronto-parietal motor circuits', *Journal of Cognitive Neuroscience*, 2005, 17(2), 273–81.

For activation of tongue muscles by listening to speech, see L. Fadiga et al., 'Speech listening specifically modulates the excitability of tongue muscles: a TMS study', *European Journal of Neuroscience*, 2002, 15(2), 399–402.

A good review of mirror neuron research is G. Buccino et al., 'Functions of the mirror neuron system: implications for neurorehabilitation', *Cognitive and Behavioural Neurology*, 2006, 19(1), 55–63.

Chapter 7: Visualization for Rehabilitation and Sports

That 122 mental practice papers had been published in 1980 but had increased to 20,000 by 2010 was found in: C. Schuster et al., 'Best practice for motor imagery: A systematic literature review

on motor imagery training elements in five different disciplines', *BMC Medicine*, 2011, 9(75), 1–35.

For the 2014 meta-analysis of stroke rehabilitation, see A.Y. Kho et al., 'Meta-analysis on the effect of mental imagery on motor recovery of the hemiplegic upper extremity function', *Australian Occupational Therapy Journal*, 2014, 61(2), 38–48.

For the university of Cincinnati study that used visual imagery as part of a study into rehabilitation after a stroke, see:

- S.J. Page et al., 'Mental practice in chronic stroke: results of a randomized, placebo-controlled trial', *Stroke*, 2007, 38(4), 1293–97.
- S.J. Page et al. 'Effects of mental practice on affected limb use and function in chronic stroke', *Archives of Physical Medicine and Rehabilitation*, 2005, 86(3), 399–402.

For the Chinese study that showed brain activation in stroke patients due to imagery training, see H. Liu et al., 'Mental practice combined with physical practice to enhance hand recovery in stroke patients', *Behavioural Neurology*, 2014, Article ID 874416.

For use of mental imagery in rehabilitation after spinal cord injury, see S.C. Cramer, et al., 'Effects of imagery training after chronic, complete spinal cord injury', *Experimental Brain Research*, 2007, 177(2), 233–42.

For use of mental imagery by Parkinson's patients, see R. Tamir et al., 'Integration of motor imagery and physical practice in group treatment applied to subjects with Parkinson's disease', *Neurorehabilitation and Neural Repair*, 2007, 21(1), 68–75.

For the golf chip shot visualization study, see M. Brouziyne and C. Molinaro, 'Mental imagery combined with physical practice

of approach shots for golf beginners', *Perceptual and Motor Skills*, 2005, 101, 203–211.

For the handball visualization study, see A. Azimkhani et al., 'The combination of mental and physical practices is better for instruction of a new skill', *Nigde University Journal of Physical Education and Sports Sciences*, 2013, 7(2), 179–187.

For the Kim studies on action observation, mental imagery and mental representation structure applied to putting in golf, see:

- T.H. Kim et al., 'Differences in learning facilitatory effect of motor imagery and action observation of golf putting', *Journal of Applied Sciences*, 2011, 11(1), 151–156.
- T. Kim, et al., 'A systematic investigation of the effect of action observation training and motor imagery training on the development of mental representation structure and skill performance', *Frontiers in Human Neuroscience*, 2017, 11(499), 1–13.

Chapter 8: Visualizing to Enhance the Immune System

For the Achterberg research that showed increases in s-IgA after immune system targeted visualization, see M. Rider, et al., 'Effect of Immune System Imagery on Secretory IgA', *Biofeedback and Self-Regulation*, 1990, 15(4), 317–333.

For the Achterberg research on cell-specific imagery of the immune system, see M. S. Rider and J. Achterberg, 'Effect of Music-Assisted Imagery on Neutrophils and Lymphocytes', *Biofeedback and Self-Regulation*, 1989, 14(3), 247–257.

For the study where 20 patients with depressed WBC counts visualized their immune systems, see V.W. Donaldson, 'A Clinical Study of Visualization on Depressed White Blood Cell Count in Medical Patients', *Applied Psychophysiology and Biofeedback*, 2000, 25(2), 117–128.

Chapter 9: Visualization for Cancer and Other Conditions

For the study of women with advanced breast cancer that found that quality of life was a predictor of their response to chemotherapy and likelihood of survival, see S.C. Fraser et al., 'A Diary of Quality of Life Measurement in Advanced Breast Cancer Trials', *British Journal of Cancer*, 1993, 67(2), 341–346.

For the 1999 study of 96 cancer patients who used visualization of their immune systems destroying cancer cells, see L.G. Walker et al., 'Psychological, Clinical, and Pathological Effects of Relaxation Training and Guided Imagery During Primary Chemotherapy', *British Journal of Cancer*, 1999, 80(1/2), 262–268.

For the RCT of 80 women who used visualization of their immune systems destroying cancer cells, see O. Eremin et al., 'Immuno-Modulatory Effects of Relaxation Training and Guided Imagery in Women with Locally Advanced Breast Cancer Undergoing Multimodality Therapy: A randomised controlled trial', *The Breast*, 2009, 18, 17–25.

For the study of women receiving surgery for breast cancer who visualized their immune systems destroying cancer cells, see C.A. Lengacher et al., 'Immune Responses to Guided Imagery During Breast Cancer Treatment', *Biological Research for Nursing*, 2008, 9(3), 205–214.

For the 1988, year-long study, of patients who used visualization of their immune systems destroying cancer cells, see B.L. Gruber et al., 'Immune System and Psychological Changes in Metastatic Cancer Patients Using Relaxation and Guided Imagery: A pilot study', *Scandinavian Journal of Behavior Therapy*, 1988, 17(1), 25–46.

For the study of 208 cancer patients who reduced cluster symptoms through the use of imagery, see A. Charalambous

et al., 'Guided Imagery and Progressive Muscle Relaxation as a Cluster of Symptoms Management Intervention in Patients Receiving Chemotherapy: A randomized controlled trial', *PLoS ONE*, 2016, 11(6), e0156911.

For the study of 65 breast cancer patients who reduced many of their side effects through the use of imagery, see S.F. Chen et al., 'Effect of Relaxation With Guided Imagery on The Physical and Psychological Symptoms of Breast Cancer Patients Undergoing Chemotherapy', *Iranian Red Crescent Medical Journal*, 2015, 17(11), e31277.

For the study where mental imagery helped wean patients from mechanical ventilation, see LeeAnna Spiva et al., 'The Effects of Guided Imagery on Patients Being Weaned from Mechanical Ventilation', *Evidence-Based Complementary and Alternative Medicine*, 2015, Article ID 802865, 1–9.

For the Hong Kong study where visualization benefitted patients with COPD, see S. W.S. Louie, 'The effects of guided imagery relaxation in people with COPD', *Occupational Therapy International*, 2004, 11(3), 145–59.

For the Alaskan study where biologically targeted imagery helped asthma sufferers, see L.W. Freeman and D. Welton, 'Effects of imagery, critical thinking, and asthma education on symptoms and mood state in adult asthma patients: a pilot study', *The Journal of Alternative and Complementary Medicine*, 2005, 11(1), 57–68.

For the use of visualization in total knee replacement surgery, see A.F. Jacobsen et al., 'Guided Imagery for Total Knee Replacement: A randomized, placebo-controlled pilot study', *The Journal of Alternative and Complementary Medicine*, 2016, 22(7), 563–575.

For the use of guided imagery for treatment of osteoarthritis in older women, see C.L. Baird and L.P. Sands, 'Effect of guided imagery with relaxation on health-related quality of life in older women with osteoarthritis', *Research in Nursing & Health*, 2006, 29(5), 442–451.

For use of guided imagery for treatment of interstitial cystitis, see D.J. Carrico et al., 'Guided imagery for women with interstitial cystitis: results of a prospective, randomized controlled pilot study', *Journal of Alternative and Complementary Medicine*, 2008, 14(1), 53–60.

For use of guided imagery in wound healing after gallbladder removal surgery (cholecystectomy), see C. Holden-Lund, 'Effects of relaxation with guided imagery on surgical stress and wound healing', *Research in Nursing & Health*, 2007, 11(4), 235–244.

For the study that examined the effect of visualization ability, see:

- E. Watanabe et al., 'Differences in relaxation by means of guided imagery in a healthy community sample', *Alternative Therapies in Health and Medicine*, 2006, 12(2), 60–66.

See also:

- E. Watanabe et al., 'Effects among healthy subjects of the duration of regularly practicing a guided imagery program', *BMC Complementary and Alternative Medicine*, 2005, 5, 21.
- K. Kwেকেboom et al., 'Imaging ability and effective use of guided imagery', *Research in Nursing and Health*, 1998, 21(3), 189–98.

For the use of guided imagery for treatment of fibromyalgia pain, see E.A. Fors et al., 'The effect of guided imagery

and amitriptyline on daily fibromyalgia pain: a prospective, randomized, controlled trial', *Journal of Psychiatric Research*, 2002, 36(3), 179–87.

For the study where volunteers imagined eating M&M's, see C.K. Morewedge et al., 'Thought for Food: Imagined consumption reduces actual consumption', *Science*, 2010, 330(6010), 1530–1533.

Chapter 11: To Stress, or Not to Stress

The reference for the 2004 meta-analysis linking stress with the immune system is S. Segerstrom and G.E. Miller, 'Psychological stress and the human immune system: a meta-analytic study of 30 years of inquiry', *Psychological Bulletin*, 2004, 130(4), 601–30.

For the effects of stress on composition of wound fluid, see E. Broadbent et al., 'Psychological stress impairs early wound repair following surgery', *Psychosomatic Medicine*, 2003, 65, 865–69.

For the effects of stress on HIV and how it increases viral replication, see S.W. Cole et al., 'Impaired response to HAART in HIV-infected individuals with high autonomic nervous system activity', *Proceedings of the National Academy of Sciences, USA*, 2001, 98(22), 12695–700.

For the 18-month study linking shyness with rate of viral replication, see S.W. Cole et al., 'Psychological risk factors for HIV pathogenesis: mediation by the autonomic nervous system', *Biological Psychiatry*, 2003, 54(12), 1444–56.

For the study where students wrote about traumatic experiences on four consecutive days, see J.W. Pennebaker and S.K. Beall, 'Confronting a traumatic event: toward an understanding of inhibition and disease', *Journal of Abnormal Psychology*, 1986, 95(3), 274–81.

For the effect of writing about traumatic experiences and the hepatitis B vaccination, see K.J. Petrie et al., 'Disclosure of trauma and immune response to hepatitis B vaccination program', *Journal of Consulting and Clinical Psychology*, 1995, 63(5), 787–92.

For the effect of writing on viral load and CD4 cell counts of HIV patients, see K.J. Petrie et al., 'Effect of written emotional expression on immune function in patients with Human Immunodeficiency Virus infection: a randomized trial', *Psychosomatic Medicine*, 2004, 66, 272–275.

For effects of emotional support on the health of breast cancer patients, see B.L. Andersen et al., 'Distress reduction from a psychological intervention contributes to improved health for cancer patients', *Brain, Behavior, and Immunity*, 2007, 21(7), 953–61.

For the effect of mindfulness-based stress reduction (MBSR) on the health of breast and prostate cancer patients, see:

- L.E. Carlson et al., 'One-year pre-post intervention follow-up of psychological, immune, endocrine and blood pressure outcomes of mindfulness-based stress reduction (MBSR) in breast and prostate cancer patients', *Brain, Behavior, and Immunity*, 2007, 21(8), 1038–49.
- L.E. Carlson et al., 'Mindfulness-based stress reduction in relation to quality of life, mood, symptoms of stress and levels of cortisol, dehydroepiandrosterone sulfate (DHEAS) and melatonin in breast and prostate cancer outpatients', *Psychoneuroendocrinology*, 2004, 29(4), 448–74.
- M. Speca et al., 'A randomized, wait-list controlled clinical trial: the effect of a mindfulness meditation-based stress reduction program on mood and symptoms of stress in cancer outpatients', *Psychosomatic Medicine*, 2000, 62(5), 613–22.

For use of MBSR in control of glucose levels in type 2 diabetes patients, see S. Rosenzweig et al., 'Mindfulness-based stress reduction is associated with improved glycemic control in type 2 diabetes mellitus: a pilot study', *Alternative Therapies in Health and Medicine*, 2007, 13(5), 36–8.

For improvement in mood, and for reduced stress and anxiety in healthy adults using meditation, see J.D. Lane et al., 'Brief meditation training can improve perceived stress and negative mood', *Alternative Therapies in Health and Medicine*, 2007, 13(1), 38–44.

For the effect of meditation at the genetic level, see J.A. Dusek et al., 'Genomic counter-stress changes induced by the Relaxation Response', *PLoS ONE*, 2008, 3(7), e2576, 1–8.